

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
13 May 2004 (13.05.2004)

PCT

(10) International Publication Number
WO 2004/039985 A2

(51) International Patent Classification⁷: C12N 15/80

MALLOY, Shannon [/GB]; Horticulture Research International, Wellesbourne, Warwick, Warwickshire, CV35 9EF (GB).

(21) International Application Number:
PCT/GB2003/004716

(74) Agent: LORD, Hilton, David; Marks & Clerk, 57-60 Lincoln's Inn Fields, London WC2A 3LS (GB).

(22) International Filing Date: 31 October 2003 (31.10.2003)

(25) Filing Language: English

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(26) Publication Language: English

(30) Priority Data:
0225390.4 31 October 2002 (31.10.2002) GB

(71) Applicant (*for all designated States except US*): HORTICULTURE RESEARCH INTERNATIONAL [/GB]; Wellesbourne, Warwick, Warwickshire CV35 9EF (GB).

(84) Designated States (*regional*): ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): BURTON, Kerry [/GB]; Horticulture Research International, Wellesbourne, Warwick, Warwickshire, CV35 9EF (GB). CHALLEN, Michael [/GB]; Horticulture Research International, Wellesbourne, Warwick, Warwickshire, CV35 9EF (GB). ELLIOTT, Timothy [/GB]; Horticulture Research International, Wellesbourne, Warwick, Warwickshire, CV35 9EF (GB). SREENIVASAPRASAD, Surapareddy [/GB]; Horticulture Research International, Wellesbourne, Warwick, Warwickshire, CV35 9EF (GB). EASTWOOD, Daniel [/GB]; Horticulture Research International, Wellesbourne, Warwick, Warwickshire, CV35 9EF (GB).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: SELECTIVE EXPRESSION IN FILAMENTOUS FUNGI

(57) Abstract: Gene promoters activated only during development of the fruiting body of filamentous fungi permit expression of large quantities of heterologous genes, even where such are otherwise toxic to the organism.

WO 2004/039985 A2

SELECTIVE EXPRESSION IN FILAMENTOUS FUNGI

The present invention relates to the transformation of filamentous fungi with heterologous DNA, methods for such transformation and processes for the collection of the resulting expression products.

Filamentous fungi have an unusual life cycle, in that the majority of the cycle is spent as mycelium, which is a largely hidden network of filaments permeating throughout and over a suitable food source. In the case of *Agaricus bisporus*, or button mushroom, which is a horticultural adaptation of a wild mushroom, the food source is decomposed leaf litter, in the wild, or generally composted cereal straw, when cultivated, and little signs of growth can be observed above ground level, other than the characteristic smell, until the fruiting bodies, or sporophores, start developing.

Compared with the culture of other vegetables, the commercial production of button mushrooms is complex. They are best grown indoors in controlled environmental conditions in trays, shelves or bags filled with a speciality compost, specifically made by, and for the, the mushroom industry from ingredients including: plant remains, such as cereal straw, corn husks, hay, rape straw; animal manure, such as from horse, chicken, cow, pig; gypsum, and other additives.

Mushroom compost is made by the mixing of the raw ingredients, for example straw, manure and gypsum base, with water, which leads to microbial degradation and heat production. The composting process has two phases. In Phase I composting, the mixture is either stacked in a windrow outdoors, or in a barn, and turned every 2 to 3 days over a three-week period, or it is placed in a walled bunker and aerated from below over an 8-10 day period. This compost is then subjected to Phase II composting in an aerated bunker where the temperature is raised to 58-60°C for approximately 12 hours and held at 50-54°C for 5 days. The compost is then ready for spawning.

Spawn is, typically, sterilised cereal grain, such as rye or millet, which has been colonised with the mushroom mycelium. Spawn is mixed with the compost and

maintained at 24°C. After approximately 17 days at this temperature, the compost is fully colonised by the mycelium.

A casing layer is then applied to the surface of the colonised compost to a depth of approximately 5 cm. Casing normally consists of peat (or peat substitute for example, coir, bark, mineral washing), chalk/lime/sugarbeetlime, water and a small amount of specialised spawn, which may be the mycelium cultured on compost, grain, vermiculite or other substrate. When the surface of the casing layer is colonised with mycelium, the temperature is lowered to 16-18°C, which induces production of mushroom sporophores, or fruiting bodies. From application of the casing to harvesting the first sporophore takes approximately 17 days.

The production of mushrooms occurs periodically, over cycles known as flushes, at 7-10 day intervals. High productivity farms normally crop mushrooms over 3 flushes. The time from the first harvest of the first flush to the last harvest of the third flush is about 21 days.

The time each growing house is occupied with each crop is from after casing and during crop production, or $17 + 21 = 38$ days, or thereabouts. This allows a growing house to be used for 9 crops per year. High productivity farms routinely have yields of 25kg/m². Because mushroom production is not light dependent, the crops are grown in layers, either four or six to maximise productivity. Accordingly, the four-layer system is expected to produce 3,200 tonnes per annum in a hectare of growing rooms.

The rapid development of the fruiting body is unique in the vegetable kingdom, and represents a significant target for the expression of suitable genes. If this rapid, volumetric expansion of the fruiting body could be harnessed together with expression of a suitable gene, then this might make possible the production, on an industrial scale, of biological products that currently defy large scale production.

Attempts to express heterologous proteins in filamentous fungi have met with numerous problems, not least being the difficulty associated with transforming the fungi in the first place. For example, WO95/02691 relates to transforming a mushroom mycelium and fruit bodies through methods such as electroporation with suitable vectors. Although this method works on a small scale, it is not especially efficient.

WO98/45455 relates to the possibility of transforming moulds, such as *Agaricus bisporus*, with the *Agrobacterium tumefaciens* bacterium, which causes crown gall tumours at the wound site of infected dicotyledonous plants. This bacterium is well known for its ability to transform plants, but it has only recently been established that it can also transform filamentous fungi.

WO 96/41882 discloses the expression of hydrophobins by the *hyp A*, *hyp B*, *hyp C* and *hyp D* genes. These are naturally occurring fungal products, and are expressed during fruiting. Heterologous expression is suggested, in connection with enhancing the flavour and/or nutritional content of the fruiting bodies.

Mol. Gen. Genet. (1993), 238: 91-96, provides a reporter-gene system in *S. commune*, for the expression of hydrophobins.

The problem still remains that the expression of heterologous genes in substantial amounts will generally substantially reduce the growth potential of the mycelium and, therefore, the harvest of the fruiting body, and prohibits the expression of any substance which is, in any way, toxic to the growth of the fungus.

Surprisingly, it has now been found that at least three genes are switched on, or otherwise subjected to elevated levels of expression, at around the veil-break stage of fruiting body development, and that heterologous DNA under the control of the expression mechanisms of these fungal genes can be selectively expressed at this stage of development of the fungus, rather than during growth of the mycelium.

Thus, in a first aspect, the present invention provides a filamentous fungus transformed with a heterologous sequence of DNA, the fungus being capable of expressing the heterologous DNA, characterised in that the heterologous DNA is under the control of a filamentous fungus transcription promoter active substantially only during stage 1, or later, of the development of the fruiting body of the fungus.

It is a particular advantage of the present invention that little or no metabolic energy need be diverted from mycelium growth, thereby maximising fruiting body mass and concomitant tissue capable of expressing the heterologous gene once it is switched on.

It is a further advantage of the present invention that little or none of the heterologous gene product is expressed during vegetative growth of the mycelium, thus enabling the production of substantially any substance capable of expression in the filamentous fungus in question, even if that substance, either alone or in combination, results in the death or stasis of the fungus. The promoters allow the synchronous switching on of the gene at a time of rapid growth and high metabolism so that, by the time any potentially toxic effects become apparent, harvestable quantities of the substance are available. Where the gene product is a regulator of mushroom growth, for example, then such considerations are not generally necessary.

It is particularly preferred to harvest the expression product of the heterologous DNA, and the product may be purified, if and as desired, by any suitable means, such as are well known in the art.

Accordingly, the present invention further provides a method for the production of a substance expressible by a DNA sequence, wherein the sequence is operably associated with a filamentous fungus transcription promoter active substantially only during stage 1, or later, of the development of the fruiting body of the fungus, the sequence and promoter being expressibly incorporated in a filamentous fungus, the fungus being cultured to fruition and the product being harvested.

At present, no less than three genes have been identified that are expressed substantially only during development of the fruiting body, and particularly during stages 4 to 7 (veil-break onwards). Without being constrained by theory, it is likely that these genes are associated with the massive water uptake required for the expansion of the fruiting body and its maturation.

The three genes, so far identified, are *abst1*, *rafe* and *mag2*. The expression product of *abst1* appears to be involved in the transport of sugars, whilst the expression product of *rafe* is a putative riboflavin aldehyde forming enzyme. The expression product of *mag2* is a so far unidentified morphogenesis associated protein.

The sequences of the three genes, and associated promoter and terminator regions, are given in the accompanying Sequence Listing.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 illustrates the stages of development of the fruiting body of a mushroom fungus, from vegetative mycelium (VM) to Stage 1 through to Stage 7 .

Figure 2A illustrates the level of expression of *abst1* during the various stages of development of the fruiting body, from vegetative mycelium (VM) to Stage 1 through to Stage 7. Figure 2B illustrates the gene expression associated with the various parts of the fruiting body at stage 4. US represents upper stipe; LS represents lower stipe; C represents cap; G represents gill; and S represents skin.

Figure 3 corresponds to Figure 2, except showing the expression of *rafe*. The stages in Figure 3A are as in Figures 1 and 2A, above. In Figure 3B, the parts of the fruiting body in which expression of *rafe* are shown, is as in Figure 2B, above.

Figure 4 corresponds to Figures 2 and 3 above, showing the expression of *mag2*. The stages and fruiting body parts showing expression are as above.

Figure 5 shows a construct with restriction enzyme sites represented by numbers 1-6.

Figure 6 shows a promoter - intron cassette, comprising *A. bisporus* *GPD* promoter 5'UTR sequence, an ATG start codon and three naturally occurring introns (*in1*, *in2* and *in3*). The restriction sites *KpnI* and *NarI* are shown. The nucleotide sequence of this cassette is given in SEQ ID NO. 12. The ATG (start codon) and VKV.....(FQ).....GRIG(R) have not been included in the complete amino acid sequence.

Figure 7 shows a leader (L) sequence, the *SPR* leader sequence, comprising 57 nucleotides (SEQ ID NO. 13) encoding 19 amino acids (SEQ ID NO. 14). This leader sequence is derived from Y13805 deposited as Accession Number Y13805 in the EMBL/GenBank/DDBJ databases (Kingsnorth, C. S., Eastwood, D. C. & Burton, K. S. (2000). *Agaricus bisporus* partial mRNA for serine proteinase).

Figure 8 shows a Terminator cassette, corresponding to Figure SEQ ID NO. 35, comprising a *A. nidulans trpC* terminator 3'UTR and a number of engineered restriction sites (*Bam* HI, *Bgl*II and *Kpn*I).

Figure 9 shows a construct in a pBluescript plasmid, combining *A. bisporus GPD* promoter with introns (as detailed above, referred to as *P*) fused with *eGFP* gene and *A. nidulans trpC* terminator (as described above, referred to as *T*). Engineered *Kpn*I sites are used to excise expression cassette from pBluescript and transfer to *Agrobacterium* binary vector harbouring hygromycin (*hph*) resistance selectable marker. T-DNA introduced into *A. bisporus* by Agro-transfection contains *hph* and heterologous protein (GFP) as a pair of divergently transcribed genes.

It will be appreciated that the Basidiomycetes, including members of the *Agaricales*, of which *A. bisporus* is one, share the exceedingly rapid development of the fruiting body in common. Without being constrained by theory, it is envisaged that this development is as a result of a rapid increase in osmotic pressure in the cells of the immature fruiting body, thereby causing a rapid influx of water into the cells. The resulting sudden expansion of the cells expands the fruiting body up to several hundred times its original size.

One or more sugar transport mechanisms are switched on at the early stages of fruiting body development, and abundant expression product is noted, especially by stage 4. These genes form a preferred subject of the present invention, and especially the control element associated therewith, but it will be appreciated that any gene selectively expressed, or with greatly enhanced expression, during development of the fruiting body is useful in the present invention.

The *abst1* gene is up-regulated, by more than 100-fold, during mushroom development, and is abundantly expressed through stages 4 to 7, and represents about 0.6% of the transcripts detected at stage 4. The transcript is about 1.7kb in length.

The gene product of *rafe* is up-regulated by about 50-fold during mushroom development, and is abundantly expressed from stage 4 onwards, as with *abst1*. The transcript is 0.7kb in length.

The expression product of *mag2* is up-regulated by about 30-fold during mushroom development, and represents about 0.2% of the transcripts at stage 4. Unlike *abst1* and *rafe*, the expression of *mag2* appears to be comparable in both the stipe and cap tissues. The length of the transcript is about 0.7kb.

The control elements, and especially the promoters, of these and other genes expressed during the development of the fruiting body are particularly useful in the present invention. Elements from genes associated with sugar transport are particularly preferred.

It is a particular advantage of the present invention that heterologous genes can be expressed at selected stages of sporophore development, where these genes might otherwise be harmful to the fungus. Expression of the heterologous genes occurs substantially only during growth of the fruiting body so that, unless the gene product is acutely toxic, then large amounts of the gene product can be expressed which would otherwise harm or hinder growth of the fungus.

Owing to the commonality of the fruiting body growth process, the promoters of any one filamentous fungus, switched on during the fruiting body growth cycle, may be employed in other filamentous fungi in the context of the present invention, in order to express heterologous genes.

In the accompanying Sequence Listing, SEQ ID NO's 12 and 13 are the promoter sequences associated with *abst1* and *rafe*, respectively, and the full sequences are provided for *abst1* and *rafe* as SEQ ID NO's 4 and 9. These include a substantial portion of the upstream promoter area and, following the procedures described in the accompanying Example, the upstream promoter area is also obtainable for *mag2*.

The promoter may be used in association with other suitable control sequences, such as terminators. A suitable terminator may be as shown in the accompanying

sequences, or may be the *Aspergillus nidulans* trpC terminator, for example. Other terminators are well known in the art.

The terminator sequences of *abst 1* and *rafe* are provided as SEQ ID NO's 35 and 36, and it has been found that it is desirable to use terminators generally associated with the promoters, so that it is preferred to use the *abst 1* terminator sequence with the *abst 1* promoter sequence, and likewise for *rafe*. However, it will be appreciated that the present invention also envisages any suitable expression system comprising the promoter, and any suitable terminator may be employed, as desired.

It will also be appreciated that the promoter sequences and terminator sequences of the invention are preferably those as listed as SEQ ID NO's 12 and 13, and 35 and 36, respectively, and sequences comprising these sequences, as well as sequences hybridising with these sequences, preferably under conditions of 60% stringency or higher, provided that promoter activity is retained in the by the sequence or the sequence to which it hybridises. Mutations and naturally occurring variants of the sequences are encompassed, and it may be, for example, appropriate to introduce a restriction site or sites for ease of manipulation. Provided that promoter activity is retained, there is no restriction on how much the promoter may be modified. Similar considerations apply to the terminator sequences.

The promoter regions may be used in their entirety when preparing heterologous genes for expression in filamentous fungi. Alternatively, it may be preferred to use consensus sequences from these regions. There is no especial advantage to using consensus sequences, except that these may be shorter. Otherwise, it is sufficient to supply the promoter upstream of the desired heterologous gene. Being a promoter, there is also no requirement that it be in the correct reading frame, just within the appropriate promoter distance.

It is generally preferable that the transformed fungus also expresses a linked selectable marker. Any marker known in the art may be used, and may be excised once a faithful strain has been generated. However, it is generally preferable that the transformed fungus maintains a marker to ensure that the desired heterologous product is still produced, and to ensure that there is no reversion to wild type. In this respect, it

is preferred that the marker have no significant negative effect on either the fungus or the product. Such markers may normally be selected from resistance markers, in order that the growth medium contain amounts of an antifungal agent ensuring that only transformed fungus can grow successfully.

Suitable markers include the hygromycin resistance cassette and the benomyl resistance tubulin gene.

Suitable methods for transforming filamentous fungi are as described above with respect to WO95/02691 and WO98/45455, which disclosures are incorporated herein by reference.

In general, the desired control sequences are ligated with the appropriate heterologous expression sequences and prepared for insertion into a suitable preparation of the fungus, such as protoplasts, all by methods well known in the art.

The resulting organism can then be grown by standard methods, and prepared as spawn after cultivation of the resulting mycelium. Spawn has the advantage that it can be stored inert for relatively long periods of up to about a year, although it is generally preferred to use it within about 4 months.

Spawn may be produced in any recognised manner, such as by growing the mycelium on sterile agar and introducing the culture to autoclaved grain. The grains may then be stored at elevated temperature to encourage colonisation, and then kept at reduced temperatures until needed.

It will be appreciated that the heterologous gene for incorporation may be in the form of cDNA or genomic DNA. cDNA is preferred, as it is generally shorter and more easy to handle.

It will also be appreciated that the heterologous gene insert should encode the sequence desired, including leader sequences and cleavage sequences, if required.

It will also be appreciated that greater expression may be achieved if fungal codons are used in place of mammalian codons, although expression will still occur, and such substitution is not necessary.

It will further be appreciated that heterologous genes may need to be expressed in the form of a cassette, for example, in order to produce the required product. In general, it is preferred to require as few heterologous gene products as possible, as the greater the number, the more likely it is that the fungal metabolism will interfere in some way, and it is generally desirable to minimise unpredictability.

Thus, although not limited thereto, it is generally preferred to limit the number of heterologous expression products to one, two or three, preferably one or two, and preferably one, other than any marker. The marker is preferably linked to the heterologous gene, such as downstream of the gene and also under the control of the fungal promoter, so as best to indicate successful and/or continuing stable transformation.

It is particularly preferred that products such as peptides be the target, as these can be harvested relatively simply. Thus, enzymes and antibodies are particularly useful, although conformational proteins, such as vaccine antigens, and active peptides such as interferons are also useful.

Accordingly, heterologous genes suitable for expression in the filamentous fungi include those whose expression results in the production of: antibodies, including other diagnostic material; secondary metabolites, such as lectins, pesticidal compounds such as *Bacillus thuringiensis* toxin (Bt toxin); therapeutic compounds such as vaccines, steroids, heterocyclic organic compounds; biological macromolecules, such as interferon, endostatin and insulin; and medical enzymes, such as thrombolytics and cerebrosidases.

In the context of the present invention, the term "heterologous" includes native DNA not normally associated with heightened expression during sporophore production. In such a respect, the native gene becomes heterologous insofar as its expression pattern is altered. Such expression may generally serve one of two purposes. The first is generally to obtain large/greater amounts of native protein, such as by transforming the filamentous fungi with extra copies or modified copies of a native gene or genes. The second may be used instead to affect/control the characteristics of mushroom crop production, such as by altering the timing of crop, flushing pattern,

yield, growth rate and/or final size of the mushroom sporophore. This latter may also suitably be achieved by the introduction of heterologous DNA from other species, if desired.

The crops are preferably allowed to go to full cap development, where possible, in order to maximise expression of the heterologous gene, although the skilled person will appreciate the best stage for harvesting any given product. The resulting caps may then be processed in any suitable manner to extract and/or purify the product, or the caps may otherwise be employed or processed, as desired.

Where the product is potentially dangerous, standard procedures may be employed between crops to entirely sterilise the area, such as steam sterilisation and swabbing of the walls, as described above.

The present invention will now be illustrated further, by reference to the following, non-limiting Examples.

EXAMPLE 1

Mushroom strains and growing conditions

A commercial *A. bisporus* strain U3 (Sylvan, U.K.) and a carboxin resistant *A. bisporus* mutant C54-*carb*.8 were used in this work. Vegetative mycelium was produced on sterile compost at 25°C and fully colonised compost (21 days) was frozen in liquid nitrogen. Mushrooms were grown in trays according to commercial practice at the Horticulture Research International mushroom cropping unit. Sporophores were produced in synchronous weekly flushes and mushroom fruit bodies from second flush were harvested at developmental stages one to seven and flash frozen. For tissue expression analysis, stage 4 mushrooms were dissected into stipe (upper and lower), cap (pileus trama), skin (pilei pellis) and gills (lamellae) and frozen. All frozen samples were stored at -80°C.

Bacterial strains, vectors and phagemid/cosmid DNA extraction

Escherichia coli strains XL-1 Blue and XL0LR (Stratagene) were used for the preparation and propagation of cDNA clones. Phagemid and Cosmid DNA extractions were carried out using the Tip 20 plasmid DNA extraction kit (Qiagen).

Preparation of an ordered library of random cDNA clones

Total RNA extraction and poly (A)⁺ RNA isolation were carried out as described previously. Using 5 µg poly (A)⁺ RNA, cDNA libraries (ZAP EXPRESS, Stratagene) were constructed from veil break stage mushrooms, and mushrooms were harvested and stored for two days. Mass excision of the cDNA libraries was performed according to the manufacturer's (Stratagene) instructions. Clear single colonies (cDNA clones in pBK-CMV) were picked and ordered in microtitre plates containing 200 µl media 96 broth with Kanamycin (50 µg ml⁻¹) in each well.

Differential screening

Duplicate nylon membranes containing DNA from the clones were prepared for differential screening. For preparing probes cDNA was generated from 5 µg poly (A)⁺ RNA using the Ready-to-Go T-primed cDNA synthesis kit (Amersham Pharmacia Biotech). Total cDNA's were labelled with [α -³²P] dCTP using the Redi-prime random labelling kit (Amersham Pharmacia Biotech). One set each of the membranes containing 3500 cDNA clones from the veil break stage mushrooms were hybridised with the cDNA probe from the veil break stage mushrooms and the cDNA probe from button stage mushrooms. Putative differentially expressed cDNA clones were re-screened to reduce false positives.

Northern analysis

Total RNA was extracted from compost colonised vegetative mycelium and seven different developmental stages of the mushroom, freshly harvested button stage mushroom and samples stored from 1 – 5 days as well as from the stipe, cap and gill tissue. Equal quantities (15 µg) of total RNA for each sample were electrophoresed and Northern blots prepared as described elsewhere. cDNA randomly labelled with [α - 32 P] dCTP, as described above, was used as the probe.

Genomic library screening and identification of the gene

For isolating cosmid clones containing the gene, a genomic library constructed from a carboxin resistant mutant of *A. bisporus* C54-*carb*.8 in cosmid vector Lawrist was used. Preliminary screening of DNA pools from 56 microtitre plates (96 clones each, *ca.* 5376) was done by PCR using primers designed from the cDNA sequence. Individual cosmid clones containing the gene were identified by probing colony blots of the 96 clones in each of the positive pools.

Nucleotide sequence determination and analysis

cDNA's in the phagemid vector pBK-CMV were initially sequenced using the vector primers T3 and T7 and the full double strand sequence was obtained using additional primers synthesised from known sequence. Genomic sequence was generated from cosmid clones by primer walking, where the initial sequencing was by done using the cDNA primers and further primers for sequencing were designed from known sequence. Nucleotide sequences were determined using the ABI automated DNA sequencing technology. Sequencing reactions were carried by thermal cycling using the ABI PrismTM BigDye terminator cycle sequencing kit (ABI-Perkin Elmer) as per manufacturers' instructions. Editing and assembling of the sequence data were done using the programmes within the DNASTAR package (Lasergene software, Dnastar

Inc.). Homology searches and nucleotide and amino acid sequence comparisons were made using a suite of software available on the WWW, particularly ExPASy (www.expasy.com) and other linked sites.

EXAMPLE 2

Constructs

Constructs are prepared to maximise expression of specific proteins through the use of homologous regulatory sequences from the mushroom *Agaricus bisporus*. The sequences comprise (see Figure 5):

- *promoter sequences P*: 5' untranslated regions preceding *A. bisporus* genes previously shown to be highly expressed in specific mushroom tissues or developmental stages;
- *leader sequences L* (optional): specific N-terminal regions of proteins that encompass the ATG nucleotide start codon for the Methionine amino acid and a defined number of other amino acids implemented in the cellular 'targeting' of proteins;
- *intron sequences I* (optional): the protein coding sequence is interrupted with a short region(s) of non-coding sequence. Intron sequences are naturally occurring in many eukaryotes and have been defined in all *A. bisporus* genes hitherto characterised. In the absence of *L* sequences the *I* cassette encompasses an ATG start codon that precedes the intron sequences;
- *gene/protein sequences G*: coding sequences for the desired heterologous protein and encompassing a stop codon;
- *terminator region T*: 3' untranslated regions following defined genes from *A. bisporus*, or other fungi, and implemented in the accurate termination and stability of gene transcripts.

P, *L*, *I*, *G* and *T* sequences are engineered from defined nucleic acids using the 'polymerase chain reaction' with defined oligonucleotide primers to introduce specific and novel restriction enzyme sites that facilitate the assembly of constructs through

molecular cloning protocols. An example construct is shown in Figure 5 with restriction enzyme sites represented by number 1-6. Therefore, cassette *P* with engineered restriction sites 1 at the 5' end and 2 at the 3' end is ligated with cassette *L* with engineered sites 2 (or compatible overhang sequence) at the 5' end and 3 at the 3' end.

P, (*L*), *I*, *G* and *T* cassettes are assembled in a plasmid vector (for example pBluescript2) and amplified at appropriate stages in *E. coli* using appropriate molecular cloning protocols.

Promoter (*P*) sequences used in heterologous protein constructs:

The *ABST1* promoter sequence is shown in SEQ ID NO. 12. The *RAFE* promoter sequence is shown in SEQ ID NO. 13.

Strategy for development of promoter - intron cassette

A. bisporus GPD promoter 5'UTR sequences ATG start codon and three naturally occurring introns (in1-3; lower case). An Engineered PCR product (486 nucleotides) with restriction sites *KpnI* and *NarI* and introns is shown in SEQ ID NO. 14 and in Figure 6.

Leader (*L*) sequence

An example of a suitable leader sequence is the *SPR* leader sequence, comprising 57 nucleotides (SEQ ID NO. 15) encoding 19 amino acids (SEQ ID NO. 16), and deposited as Genbank (NCBI) Accession Number Y13805. The sequence alignment of the nucleotide and amino acids is given in Figure 7.

Intron (*I*) sequences:

In addition to *GPD* introns referred to above, numerous *A. bisporus* introns have been characterised that can be used as intron cassettes (see tables 2 to 5, below). These introns have been established by comparison of partial cDNA and genomic DNA

sequences. The introns are described by their gene of origin (i.e. putative gene product) and Genbank (NCBI) cDNA accession number. For each gene, introns are numbered from the 5' end. The first identified introns of each gene are theoretical and unconfirmed (cDNA sequence not available at 5' end and therefore identified by comparison of gene sequence with sequences of similar genes from other organisms) and are not included in this list.

Terminator (*T*) sequences:

The *abst1* terminator sequence is given in SEQ ID NO. 35. The *rafe* terminator sequence is given in SEQ ID NO. 36.

Strategy for development of terminator (*T*) cassette

A. nidulans trpC terminator 3'UTR as an engineered PCR product (comprising 800 nucleotides) with introduced restriction sites *Bgl*II and *Kpn*I, is given in SEQ ID NO. 37, and shown diagrammatically in Figure 8.

Construct strategy for expression of GFP protein:

Construct in pBluescript plasmid combines *A. bisporus GPD* promoter with introns (as detailed above; *P*) fused with *eGFP* gene and *A. nidulans trpC* terminator (as described above; *T*), and represented in Figure 9. Engineered *Kpn*I sites are used to excise expression cassette from pBluescript and transfer to *Agrobacterium* binary vector harbouring hygromycin (*hph*) resistance selectable marker. T-DNA introduced into *A. bisporus* by Agro-transfection contains *hph* and heterologous protein (GFP) as a pair of divergently transcribed genes.

In the accompanying Sequence Listing, it will be noted as follows:

Organism:	<i>Agaricus bisporus</i>
Gene designations (tentative):	<i>abst1</i> (sugar transporter); <i>rafe</i> (putative riboflavin aldehyde forming enzyme gene); <i>mag2</i> (unidentified morphogenesis associated gene)
Gene identification:	By differential screening of a cDNA library (3500 random clones) from stage 4 mushrooms
Genomic clones:	From a genomic library of <i>A. bisporus</i> strain C54- <i>carb8</i> in cosmid vector Lawrist (ca. 30 – 40 kb genomic fragments)
<i>abst1</i> :	Up-regulated (more than 100 fold) during mushroom development, abundantly expressed through stages 4 – 7 (later stages of mushroom development), ca. 0.6% transcripts at stage 4, represented by 20 clones in the differential library, 1.7 kb transcript
<i>rafe</i> :	Up-regulated (up to 50 fold) during mushroom development, abundantly expressed through stages 4 – 7, 0.7 kb transcript
<i>mag2</i> :	Up-regulated (up to 30 fold) during mushroom development, ca. 0.6% transcripts at stage 4, represented by seven clones in the differential library, comparable levels of expression in stipe and cap tissue, ca. 0.7 kb transcript

The following Table provides an index of the accompanying SEQ ID NO's, and gives an indication as to their identity. The sequence provided in SEQ ID NO 4 was obtained on the basis of single strand analysis and is for guidance only, as there are some errors and unknown nucleotides. The promoter sequence of SEQ ID NO. 12 is preferred, as it is shorter, and does not comprise as many restriction sites. In addition, it was sequenced as a double strand.

Table 1

Sequence description	SEQ ID NO
<i>Abst1</i> cDNA sequence	1
Abst1 protein encoded by ORF of SEQ ID NO 1	2
<i>abst1</i> Genomic sequence	3
<i>abst1</i> Promoter region (approx. 1000 bases upstream of ORF).	4
<i>abst1</i> Terminator region (approx. 700 bases downstream of ORF)	5
<i>rafe</i> cDNA sequence	6
Rafe protein encoded by ORF of SEQ ID NO2	7
<i>rafe</i> Genomic Sequence	8
<i>rafe</i> Promoter region (approx. 1000 bases upstream of ORF)	9
<i>rafe</i> Terminator region (approx. 1200 bases downstream of ORF):	10
<i>. mag2</i> : Partial cDNA sequence	11
<i>abst 1</i> promoter sequence	12
<i>Rafe</i> promoter sequence	13
Engineered PCR product (486 nt) with restriction sites <i>KpnI</i> and <i>NarI</i> comprising <i>A. bisporus GPD</i> promoter, 5'UTR sequences, ATG start codon and three naturally occurring introns.	14
<i>SPR</i> leader sequence (nucleotide sequence)	15
<i>SPR</i> leader sequence (amino acid sequence)	16
<i>abst 1</i> terminator sequence	35
<i>rafe</i> terminator sequence	36
<i>A. nidulans trpC</i> terminator 3'UTR as engineered PCR product (800 nt) with introduced restriction sites <i>BglII</i> and <i>KpnI</i>	37

Table 2: ASL-Argininosuccinate lyase (Genbank Accession no. AJ271691)

Intron	SEQ ID NO.
ASL intron 3	17
ASL intron 4	18
ASL intron 5	19
ASL intron 6	20

Table 3: CYP1-Cytochrome P450 I (Genbank Accession no. AJ271707)

Intron	SEQ ID NO.
CYP1 intron 6	21
CYP1 intron 7	22
CYP1 intron 8	23
CYP1 intron 9	24
CYP1 intron 10	25

Table 4: SOD-Superoxide Dismutase (Genbank Accession no. AJ271694)

Intron	SEQ ID NO.
SOD intron 2	26
SOD intron	27

Table 5: LAP-Leucine Aminopeptidase (Genbank Accession no. AJ271690)

Intron	SEQ ID NO.
LAP intron 3	28
LAP intron 4	29
LAP intron 5	30
LAP intron 6	31
LAP intron 7	32

Table 6: MET-Metallothionein (Genbank Accession no. AJ271695)

Intron	SEQ ID NO.
MET intron 2	33
MET intron 3	34

CLAIMS:

1. A filamentous fungus transformed with a heterologous sequence of DNA, the fungus being capable of expressing the heterologous DNA, characterised in that the heterologous DNA is under the control of a filamentous fungus transcription promoter active substantially only during stage 1, or later, of the development of the fruiting body of the fungus.
2. A fungus according to claim 1, which is *A. bisporus*.
3. A fungus according to claim 1 or 2, wherein the promoter is that of *abst1*.
4. A fungus according to claim 1 or 2, wherein the promoter is that of *rafe*.
5. A fungus according to claim 1 or 2, wherein the promoter is that of *mag2*.
6. A fungus according to claim 1 or 2, wherein the promoter comprises the sequence of SEQ ID NO 12, or a mutation or variant thereof, or a sequence which hybridises thereto under conditions of at least 60% stringency.
7. A fungus according to claim 1 or 2, wherein the promoter comprises the sequence of SEQ ID NO 13, or a mutation or variant thereof, or a sequence which hybridises thereto under conditions of at least 60% stringency.
8. A fungus according to any preceding claim, wherein the DNA is operably linked with a terminator comprising the sequence of SEQ ID NO 35, or a mutation or variant thereof, or a sequence which hybridises thereto under conditions of at least 60% stringency.
9. A fungus according to any preceding claim, wherein the DNA is operably linked with a terminator comprising the sequence of SEQ ID NO 36, or a mutation or variant thereof, or a sequence which hybridises thereto under conditions of at least 60% stringency.

10. A fungus according to claim 1, wherein the DNA is operably linked with a promoter comprising the sequence of SEQ ID NO. 12 and a terminator comprising the sequence of SEQ ID NO 35, or a mutation or variant of either, or a sequence which hybridises thereto under conditions of at least 60% stringency.

11. A fungus according to claim 1, wherein the DNA is operably linked with a promoter comprising the sequence of SEQ ID NO. 13 and a terminator comprising the sequence of SEQ ID NO 36, or a mutation or variant of either, or a sequence which hybridises thereto under conditions of at least 60% stringency.

12. A fungus according to any preceding claim, wherein a selectable marker is linked with the heterologous DNA.

13. A fungus according to any preceding claim, wherein the heterologous DNA is native DNA.

14. A fungus according to any preceding claim, wherein the heterologous DNA is selected such as to affect characteristics of mushroom crop production.

15. A fungus according to any preceding claim, wherein the heterologous DNA encodes: antibodies, including other diagnostic material; secondary metabolites, such as lectins, pesticidal compounds such as *Bacillus thuringiensis* toxin (Bt toxin); therapeutic compounds such as vaccines, steroids, heterocyclic organic compounds; biological macromolecules, such as interferon, endostatin and insulin; and medical enzymes, such as thrombolytics and cerebrosidases.

16. A method for the production of a substance expressible by a DNA sequence, wherein the sequence is operably associated with a filamentous fungus transcription promoter active substantially only during stage 1, or later, of the development of the fruiting body of the fungus, the sequence and promoter being expressibly incorporated in a filamentous fungus, the fungus being cultured to fruition and the product being harvested.

17. A method for the production of a substance expressible by a DNA sequence, wherein the sequence is operably associated with a filamentous fungus transcription promoter active substantially only during stage 1, or later, of the development of the fruiting body of the fungus, the sequence and promoter being expressibly incorporated in a filamentous fungus, the fungus being cultured to fruition and the product being harvested, and wherein the DNA and/or fungus is as defined in any of claims 1 to 15.

1/3





Fig.2A




Fig.2B

2/3

VM 1 2 3 4 5 6 7



Fig.3A




Fig.3B

VM 1 2 3 4 5 6 7



Fig.4A




Fig.4B




Fig.5




Fig.6

atgcatttct ctttgtcttt tgccaccctt gctctcttag tcgcttcggc tgttggt
M H F S L S F A T L A L L V A S A V G

Fig.7




Fig.8




Fig.9

SEQUENCE LISTING

<110> Horticulture Research International

<120> SELECTIVE EXPRESSION IN FILAMENTOUS FUNGI

<130> WPP86709

<150> UK 0225390.4

<151> 2002-10-31

<160> 37

<170> PatentIn version 3.1

<210> 1

<211> 1736

<212> DNA

<213> Agaricus bisporus

<220>

<221> CDS

<222> (19)..(1674)

<223>

```

<400> 1
ggctgagctc tattcatc atg gcg tcg gaa cga cag att gaa gaa ctt ccc      51
                    Met Ala Ser Glu Arg Gln Ile Glu Glu Leu Pro
                    1          5          10

agt tat aag tac gcc tat att ttg acg gca tcg gct tgc ttg gga agt      99
Ser Tyr Lys Tyr Ala Tyr Ile Leu Thr Ala Ser Ala Cys Leu Gly Ser
          15          20          25

gtg ttt cac ggg tgg gat gta ggc ctt ata gga ggc ata ctc tcg cta      147
Val Phe His Gly Trp Asp Val Gly Leu Ile Gly Gly Ile Leu Ser Leu
          30          35          40

cgg tca ttt caa gaa tat ctc ggg atc aat aca aaa aat gcc gtc aag      195
Arg Ser Phe Gln Glu Tyr Leu Gly Ile Asn Thr Lys Asn Ala Val Lys
          45          50          55

aaa gcg att cta gac gga aac atc atc tct gtg ctc caa gcc gga tgt      243

```

Lys 60	Ala	Ile	Leu	Asp 65	Gly	Asn	Ile	Ile	Ser 70	Val	Leu	Gln	Ala	Gly	Cys 75	
ttt	ttt	ggc	gcg	ctt	gga	acc	gga	tat	ctc	tct	agt	cga	ttc	ggc	cga	291
Phe	Phe	Gly	Ala	Leu 80	Gly	Thr	Gly	Tyr	Leu 85	Ser	Ser	Arg	Phe	Gly 90	Arg	
aga	ccc	tgt	ctt	att	gca	tct	ggt	att	gtg	tat	ata	act	ggc	ggt	ttg	339
Arg	Pro	Cys	Leu 95	Ile	Ala	Ser	Gly	Ile 100	Val	Tyr	Ile	Thr	Gly 105	Gly	Leu	
ctg	caa	tgc	act	gtc	ggt	ttg	gga	ccc	tcg	caa	gct	gct	gct	cta	cac	387
Leu	Gln 110	Cys	Thr	Val	Gly	Leu	Gly 115	Pro	Ser	Gln	Ala	Ala	Ala	Leu	His	
gtg	ttc	tat	att	ggc	agg	ttc	att	tct	ggt	atc	ggt	gtt	ggg	atg	gtg	435
Val	Phe	Tyr	Ile	Gly	Arg	Phe 130	Ile	Ser	Gly	Ile	Gly 135	Val	Gly	Met	Val	
tcc	act	ctc	gtg	cct	ttg	tat	att	tcg	gag	tgt	gtc	cct	agg	act	ata	483
Ser	Thr	Leu	Val	Pro	Leu	Tyr 145	Ile	Ser	Glu	Cys	Val 150	Pro	Arg	Thr	Ile 155	
cgc	ggg	cgc	tgt	act	gga	aca	ctc	caa	ttt	gcg	act	aac	agt	ggt	ctg	531
Arg	Gly	Arg	Cys	Thr 160	Gly	Thr	Leu	Gln	Phe 165	Ala	Thr	Asn	Ser	Gly 170	Leu	
atg	ctg	ggc	ttc	tgg	gtc	aac	tac	agc	gtg	tcg	aaa	aac	gtg	ccc	ttt	579
Met	Leu	Gly	Phe 175	Trp	Val	Asn	Tyr 180	Ser	Val	Ser	Lys	Asn 185	Val	Pro	Phe	
ggt	gaa	atg	caa	tgg	cga	att	ccg	tta	att	atc	caa	atg	att	ccg	agc	627
Gly	Glu	Met 190	Gln	Trp	Arg	Ile 195	Pro	Leu	Ile	Ile	Gln	Met 200	Ile	Pro	Ser	
ctc	ttg	ttc	atc	ata	gcc	atg	ttt	ttc	caa	cca	gaa	tcg	ccg	aga	tgg	675
Leu	Leu	Phe 205	Ile	Ile	Ala	Met 210	Phe	Phe	Gln	Pro	Glu	Ser 215	Pro	Arg	Trp	
ctt	gtt	gaa	cac	ggg	aaa	cac	aag	gaa	gct	gcg	acg	gta	ctg	gcg	cgt	723
Leu	Val	Glu	His	Gly 225	Lys	His	Lys	Glu	Ala	Ala	Thr 230	Val	Leu	Ala 235	Arg	
act	ggc	ggc	aag	gat	ggt	gat	cat	cct	agt	ggt	gta	cag	aca	ctg	gag	771
Thr	Gly	Gly	Lys 240	Asp	Val	Asp	His	Pro	Ser 245	Val	Val	Gln	Thr	Leu 250	Glu	
gag	atc	aag	caa	gaa	ttt	gtg	gcg	agt	aaa	caa	cca	tcg	ttt	tta	aag	819
Glu	Ile	Lys	Gln 255	Glu	Phe	Val	Ala	Ser 260	Lys	Gln	Pro	Ser	Phe 265	Leu	Lys	
cag	att	cgc	ctg	gtc	ggt	gaa	tcg	agg	gct	ggt	gcc	ctg	agg	tgc	ttt	867
Gln	Ile	Arg	Leu 270	Val	Gly	Glu	Ser 275	Arg	Ala	Val	Ala	Leu 280	Arg	Cys	Phe	
ata	cca	ccg	cta	gtg	atg	ttc	ttc	cag	cag	tgg	acg	ggt	aca	aat	gcc	915
Ile	Pro	Pro	Leu 285	Val	Met	Phe 290	Phe	Gln	Gln	Trp	Thr 295	Gly	Thr	Asn	Ala	
atc	aac	ctt	tat	agt	ccc	gaa	gta	ttc	cgt	cat	ctt	gga	atc	cat	ggc	963
Ile	Asn	Leu	Tyr	Ser 305	Pro	Glu	Val	Phe	Arg	His 310	Leu	Gly	Ile	His	Gly 315	
acc	agc	ggg	gct	ctc	ttc	gct	act	ggt	ggt	tat	ggc	gtg	gtg	aag	gtt	1011
Thr	Ser	Gly	Ala 320	Leu	Phe	Ala	Thr	Gly 325	Val	Tyr	Gly	Val	Val	Lys 330	Val	
gtt	tca	gtt	gca	ctt	gcc	ctc	act	ttt	gct	gtc	gaa	cgc	ttt	gga	cgc	1059

Val Ser Val Ala Leu Ala Leu Thr Phe Ala Val Glu Arg Phe Gly Arg	
335 340 345	
aag aga ggg ttg att ttt ggt ggt atc ggc caa gca ctt atg atg ttt	1107
Lys Arg Gly Leu Ile Phe Gly Gly Ile Gly Gln Ala Leu Met Met Phe	
350 355 360	
tgg ttg gga ggt tat agt gcc acc cac caa gac ggt act gtc agt cct	1155
Trp Leu Gly Gly Tyr Ser Ala Thr His Gln Asp Gly Thr Val Ser Pro	
365 370 375	
gcg agt cat gtt tcc att gtt gca ctc tac ttg tat ggt gca ttc ttc	1203
Ala Ser His Val Ser Ile Val Ala Leu Tyr Leu Tyr Gly Ala Phe Phe	
380 385 390 395	
tcc atg gga tgg ggc cca tta cca tgg gtc gtc gct gga gag gtt gca	1251
Ser Met Gly Trp Pro Leu Pro Trp Val Val Ala Gly Glu Val Ala	
400 405 410	
cct aac cat gtc cgc tcc ttc gcc ctc tcc atc gcc gtt gga act cat	1299
Pro Asn His Val Arg Ser Phe Ala Leu Ser Ile Ala Val Gly Thr His	
415 420 425	
tgg ctc ttc ggg ttt gtg ata tca aaa gtg acg cca att atg ttg gac	1347
Trp Leu Phe Gly Phe Val Ile Ser Lys Val Thr Pro Ile Met Leu Asp	
430 435 440	
cgt atc aaa tat ggc aca ttc cta ctc ttc gga ttc tgt tgc atg ata	1395
Arg Ile Lys Tyr Gly Thr Phe Leu Leu Phe Gly Phe Cys Cys Met Ile	
445 450 455	
gta gcg aca tgg gct tat ttc tgt cta cct gag aca agt ggg ttc gct	1443
Val Ala Thr Trp Ala Tyr Phe Cys Leu Pro Glu Thr Ser Gly Phe Ala	
460 465 470 475	
ctg gag gac atc aaa tat ctg ttc gag cga gac gtc atc att cgt tca	1491
Leu Glu Asp Ile Lys Tyr Leu Phe Glu Arg Asp Val Ile Ile Arg Ser	
480 485 490	
ttg cag gac gct ccc ggt gga aaa ata ttc ttg ggg ggg agg cgt gtg	1539
Leu Gln Asp Ala Pro Gly Gly Lys Ile Phe Leu Gly Gly Arg Arg Val	
495 500 505	
gaa tct gta gct tcg ttg aaa gag agg cgc gtt gga gtc gct ggt gag	1587
Glu Ser Val Ala Ser Leu Lys Glu Arg Arg Val Gly Val Ala Gly Glu	
510 515 520	
cag ggt gag aag ata act ggt cta aat tcg gaa ttg gaa gat gtt tcc	1635
Gln Gly Glu Lys Ile Thr Gly Leu Asn Ser Glu Leu Glu Asp Val Ser	
525 530 535	
tca aaa aaa tca aca ttg aag gaa act tca tcc gtt tga tatatagtct	1684
Ser Lys Lys Ser Thr Leu Lys Glu Thr Ser Ser Val	
540 545 550	
cctaaattcta ttgtaatgcc attttcccaa ttcaaaaaaa aaaaaaaaaa aa	1736

<210> 2

<211> 551

<212> PRT

<213> Agaricus bisporus

<400> 2

```

Met Ala Ser Glu Arg Gln Ile Glu Glu Leu Pro Ser Tyr Lys Tyr Ala
1          5          10          15

Tyr Ile Leu Thr Ala Ser Ala Cys Leu Gly Ser Val Phe His Gly Trp
20          25          30

Asp Val Gly Leu Ile Gly Gly Ile Leu Ser Leu Arg Ser Phe Gln Glu
35          40          45

Tyr Leu Gly Ile Asn Thr Lys Asn Ala Val Lys Lys Ala Ile Leu Asp
50          55          60

Gly Asn Ile Ile Ser Val Leu Gln Ala Gly Cys Phe Phe Gly Ala Leu
65          70          75          80

Gly Thr Gly Tyr Leu Ser Ser Arg Phe Gly Arg Arg Pro Cys Leu Ile
85          90          95

Ala Ser Gly Ile Val Tyr Ile Thr Gly Gly Leu Leu Gln Cys Thr Val
100         105         110

Gly Leu Gly Pro Ser Gln Ala Ala Ala Leu His Val Phe Tyr Ile Gly
115         120         125

Arg Phe Ile Ser Gly Ile Gly Val Gly Met Val Ser Thr Leu Val Pro
130         135         140

Leu Tyr Ile Ser Glu Cys Val Pro Arg Thr Ile Arg Gly Arg Cys Thr
145         150         155         160

Gly Thr Leu Gln Phe Ala Thr Asn Ser Gly Leu Met Leu Gly Phe Trp
165         170         175

Val Asn Tyr Ser Val Ser Lys Asn Val Pro Phe Gly Glu Met Gln Trp
180         185         190

Arg Ile Pro Leu Ile Ile Gln Met Ile Pro Ser Leu Leu Phe Ile Ile
195         200         205

Ala Met Phe Phe Gln Pro Glu Ser Pro Arg Trp Leu Val Glu His Gly
210         215         220

Lys His Lys Glu Ala Ala Thr Val Leu Ala Arg Thr Gly Gly Lys Asp
225         230         235         240

Val Asp His Pro Ser Val Val Gln Thr Leu Glu Glu Ile Lys Gln Glu
245         250         255

Phe Val Ala Ser Lys Gln Pro Ser Phe Leu Lys Gln Ile Arg Leu Val
260         265         270

```

Gly Glu Ser Arg Ala Val Ala Leu Arg Cys Phe Ile Pro Pro Leu Val
 275 280 285

Met Phe Phe Gln Gln Trp Thr Gly Thr Asn Ala Ile Asn Leu Tyr Ser
 290 295 300

Pro Glu Val Phe Arg His Leu Gly Ile His Gly Thr Ser Gly Ala Leu
 305 310 315 320

Phe Ala Thr Gly Val Tyr Gly Val Val Lys Val Val Ser Val Ala Leu
 325 330 335

Ala Leu Thr Phe Ala Val Glu Arg Phe Gly Arg Lys Arg Gly Leu Ile
 340 345 350

Phe Gly Gly Ile Gly Gln Ala Leu Met Met Phe Trp Leu Gly Gly Tyr
 355 360 365

Ser Ala Thr His Gln Asp Gly Thr Val Ser Pro Ala Ser His Val Ser
 370 375 380

Ile Val Ala Leu Tyr Leu Tyr Gly Ala Phe Phe Ser Met Gly Trp Gly
 385 390 395 400

Pro Leu Pro Trp Val Val Ala Gly Glu Val Ala Pro Asn His Val Arg
 405 410 415

Ser Phe Ala Leu Ser Ile Ala Val Gly Thr His Trp Leu Phe Gly Phe
 420 425 430

Val Ile Ser Lys Val Thr Pro Ile Met Leu Asp Arg Ile Lys Tyr Gly
 435 440 445

Thr Phe Leu Leu Phe Gly Phe Cys Cys Met Ile Val Ala Thr Trp Ala
 450 455 460

Tyr Phe Cys Leu Pro Glu Thr Ser Gly Phe Ala Leu Glu Asp Ile Lys
 465 470 475 480

Tyr Leu Phe Glu Arg Asp Val Ile Ile Arg Ser Leu Gln Asp Ala Pro
 485 490 495

Gly Gly Lys Ile Phe Leu Gly Gly Arg Arg Val Glu Ser Val Ala Ser
 500 505 510

Leu Lys Glu Arg Arg Val Gly Val Ala Gly Glu Gln Gly Glu Lys Ile
 515 520 525

Thr Gly Leu Asn Ser Glu Leu Glu Asp Val Ser Ser Lys Lys Ser Thr
 530 535 540

Leu Lys Glu Thr Ser Ser Val
545 550

<210> 3

<211> 3638

<212> DNA

<213> Agaricus bisporus

<220>

<221> misc_feature

<222> (3)..(3)

<223> n is an unknown nucleotide

<220>

<221> misc_feature

<222> (14)..(14)

<223> n is an unknown nucleotide

<220>

<221> misc_feature

<222> (510)..(510)

<223> n is an unknown nucleotide

<400> 3

gtncgatggg ttcntctggg ttaagttgca cgacgctttc ctttttcttt tatggcctgt	60
ctgccctttt aacgctttat ctttcggcag ccatggatgt ccttcgtcac cgtattatca	120
ctcttaatcg tgggtgtgag cacatggaaa gggtcattat cgtcatccca tgacgcggtg	180
caaaattcgt cattcagagt ggaaccgata ctaggagagg attttgaaaa ggctatcgta	240
tcttcgcatc cattcactcc aacgacatct atatcgggtt ctcaagctac gataaccgtt	300
cctcacgcat cacattcttc tctgattgta atgaccgaag accgcggttc caccctaata	360
attcgaagag cttgttgcca atgtcaaaaa aggcttttgg gcaaaccggg atggcttata	420
gagcctccag ccccagcagc taacatcggg cagagaagga aaattcatcg ggcggttgaa	480
ttatcaccgt ttggttcctg agtcatctgn ragatgtacg cagatggtga taccgtgttt	540
gattggcgcc gttggagaag aactatatta ttcgatggat tttttgttcg agtttgacac	600
agagacagag atgatagagg tttgctattg atgtagcaaa ggatcatttg acgatggcgc	660

atagggcgat ggttatcttt atgtctggaa ttataatatg tattgttccc cacttttctt 720
ttatatattat taataactaat tggaagtttc agttgttgga tgagcaaagt tgggtgcagat 780
agaaactaga attcggattc ccatatctga ggtacctttt ccttccgctg gcaatcctgg 840
ccacttcgac gtggtgacgc agagggcgcg tgctattgtt agcacatgcc atawggatcg 900
acgttgcctc tcgtacttcg cgcctaggct cgctcatgcc tcgatgcac ttttcaattc 960
gggcggttgcg tctcccaggt gcctgttaaa agggcgaaact ttagtgtaat tgtactaaca 1020
cagtcctcgc ggctgagctc tattcatcat ggcgtcgga cgacagattg aagaacttcc 1080
cagttataag tacgcctata ttttgacggc atcggcttgc ttgggaagtg tgtttcacgg 1140
gtgggatgtg caagtatcct ttwgacgctg gctcatccct tgtttgctga tggttaatgt 1200
gtagaggcct tataggaggc atactctcgc tacggtcatt tcaagaatat ctcgggatca 1260
atacaaaaaa tgccgtcaag aaagcgattc tagacgaaa catcatctct gtgctccaag 1320
ccggatgttt tgtaagtctg agcactcgtt cgaccagctt actttctttt actaacaacg 1380
tgttgaccta gttttggcgc gcttgaacc ggatatctct ctagtcgatt cggccgaaga 1440
ccctgtctta ttgcatctgg tatttgttat ataactggcg gtttgctgca atgactgtc 1500
ggtttgggac cctcgcaagc tgctgctcta cacgtgttct atattggcag gttcatttct 1560
ggtatcggtg ttgggatggt gtccactctc gtgcctttgt atatttcgga gtgtgtccct 1620
aggactatac gcgggcgctg tactggaaca ctccaatttg cgactaacag tggctctgatg 1680
ctgggctgta agtgtacctt ttgttatgct ccggggacga tactaaaagt agtctgtagt 1740
ctgggtcaac tacagcgtgt cgaaaaacgt gcccttttgt gaaatgcaat ggcgaattcc 1800
gttaattatc cagtacgtta ttggatgcaa agtgaagcat atgctgaaat tagacttgtg 1860
tcgtactttc cagaatgatt ccgagcctct tgttcatcat agccatgttt ttccaaccag 1920
aatcgccgag atggcttgtt gaacacggga aacacaagga agctgcgacg gtactggcgc 1980
gtactggcgg caaggatgtt gatcatccta gtgttgtaaca gacactggag gagatcaagc 2040
aagaatttgt ggcgagtaaa caaccatcgt ttttaaagca gattcgctg gtcggtgaat 2100
cgagggtgtg tgccytgagg wgctttatac caccgctagt gatgttcttc cagcagtgga 2160
cgggtacaaa tgccatcaac ctttatagtc ccgaagtatt ccgtcatctt ggaatccatg 2220
gcaycagckr kgmwstcttc sctamtsgtg tttatkmgmt ggtrmaggtt gtttcagttg 2280
cacttgcctc cacttttgcg gtcgaacgct ttggacgcaa gagagggttg atttttggtg 2340
gtatcggcca agcacttatg atgttttggt tgggagggtta tagtgccacc caccaagacg 2400
gtactgtcag tcctgcgagt catgtttcca ttgttgact ctacttgtat ggtgcattct 2460
tctccactga agaattggtc ctgttabcat gggctcgtcg tggagagggt gcacctaac 2520
atgtccgctc cttcgccctc tccatcgccg ttggaactca ttggctcttc gggtttgtga 2580
tatcaaaagt gacgccaatt atgttggacc gtatcaaata tggcacattc ctactcttcg 2640
gattctgttg catgatagta gcgacatggg cttatttctg tctacctgag acaagtgggt 2700

```

tcgctctgga ggacatcaaa tatctgttcg agcgagacgt catcattcgt tcattgcagg 2760
acgctccccg tggaaaaata ttcttggggg ggaggcgtgt ggaatctgta gcttcgttga 2820
aagagaggcg cgttgagtc gctggtgagc agggtgagaa gataactggt ctaaattcgg 2880
aattggaaga tgtttcctca aaaaaatcaa cattgaagga aacttcattcc gtttgatata 2940
tagtctccaa attctattgt aatgccattt tccaattca aaaggacccg ctctcgaacc 3000
gggtcagatg caattttggt cagcaatggt ttatgttgtt tccccgtaag tatgcactag 3060
aagagaacaa aacgtcacta tttgctcaat gcaggatgca cctggcgaga taatattctt 3120
gcggtgaagt cgaacaacgt ctgtagtcct gtaaaaatat acagtgagta gagggatgat 3180
gccgatgtgg aaggaagcaa ccgattacga ttcggatttg gccatgagac ggccgctctt 3240
gaacagacca acaatatccc tttaaattta atacagaatt actcaatatg cttccaagta 3300
tttcgattcc tcgaattccg tctggccgcg gcatgagcat ggacaggcgg acagaagagg 3360
ctatcgttgt attgcttcat caagcgaccc tgactagtga cttcaggcat gatcatgcgc 3420
ttagcaatct gtcccttcaa gtcgagtcct cgaattcaac agctttcaac aagtcgtgat 3480
tatttgaccc ccgactggaa tcaaattggc tcttcaaatt tcaaaccctc aatgcttcat 3540
gcttcatgcg tcatgacgca agctgtcaat tttcattttc cagttcggtc ccattctcac 3600
tctcgcccct cstaatgtct tccagaaaat ggcttccc 3638

```

<210> 4

<211> 1033

<212> DNA

<213> Agaricus bisporus

<220>

<221> misc_feature

<222> (495)..(495)

<223> n is an unknown nucleotide

<400> 4

```

ctgggttaag ttgcacgacg ctttcctttt tcttttatgg cctgtctgcc cttttaacgc 60
tttatctttc ggcagccatg gatgtccttc gtcaccgtat tatcactctt aatcgtggtg 120
gtgagcacaat ggaaagggtc attatcgtca tcccatgacg cggtgcaaaa ttcgtcattc 180
agagtgaac cgatactagg agaggatttt gaaaaggcta tcgtatcttc cgatccattc 240
actccaacga catctatatc gggttctcaa gctacgataa ccgttcctca cgcacacat 300
tcttctctga ttgtaatgac cgaagaccgc ggttccaccc ctaatatctg aagagcttgt 360
tgccaatgtc aaaaaaggct tttgggcaaa ccgggatggc ttatcgagcc tccagcccca 420

```

```

gcagctaaca tcgggcagag aaggaaaatt catcgggcgg ttgaattatc accgtttggt 480
tcctgagtca tctgnragat gtacgcagat ggtgataccg tgtttgattg gcgccgttgg 540
agaagaacta tattattcga tggatttttt gttcagagtt gacacagaga cagagatgat 600
agagggtttgc tattgatgta gcaaaggatc atttgacgat ggcgcatagg gcgatggtta 660
tctttatgtc tggaattata atatgtattg ttccccactt ttcttttata tttattaata 720
ctaattggaa gtttcagttg ttggatgagc aaagttggtg cagatagaaa ctagaattcg 780
gattcccata tctgaggtag cttttccttc cgctggcaat cctggccact tcgacgtggt 840
gacgcagagg gcgcgtgcta ttgtagcac atgccatawg gatcgacgtt gcctctcgta 900
cttcgcgcct aggtcgcctc atgcctcgat gcactttttc aattcggggc ttgcgtctcc 960
cagggtgcctg ttaaaagggc gaactttagt gtaattgtac taacacagtc cctcgggctg 1020
agctctattc atc 1033

```

<210> 5

<211> 702

<212> DNA

<213> *Agaricus bisporus*

```

<400> 5
tatatagtct ccaaattcta ttgtaatgcc attttcccaa ttcaaaagga cccgctctcg 60
aaccgggtca gatgcaatth ttgtcagcaa tggtttatgt tgtttccccg taagtatgca 120
ctagaagaga acaaaacgtc actatttget caatgcagga tgcacctggc gagataatat 180
tcttgcggtg aagtcgaaca acgtctgtag tcctgtaaaa atatacagtg agtagaggga 240
tgatgccgat gtggaaggaa gcaaccgatt acgattcgga tttggccatg agacggccgc 300
tcttgaacag accaacaata tccctttaa tttaatacag aattactcaa tatgettcca 360
agtatttcga ttctcgaat tccgtctggc cgcggcatga gcatggacag gcggacagaa 420
gaggctatcg ttgtattget tcatcaagcg accctgacta gtgacttcag gcatgatcat 480
gcgcttagca atctgtccct tcaagtcgag tccccgaatt caacagcttt caacaagtcg 540
tgattatttg acccccgact ggaatcaaat tggtcttca aatttcaaac cttcaatget 600
tcatgcttca tgcgtcatga cgcaagctgt caattttcat tttccagttc ggtcccatc 660
tcactctcgc ccctcstaat gtcttcaga aaatggcttc cc 702

```

<210> 6

<211> 689

<212> DNA

<213> *Agaricus bisporus*

<220>

<221> CDS

<222> (147)..(521)

<223>

<400> 6

```

cgcctacttt tataccaacc ccaaattccaa aggttgaaaa aaaaatttcg acaaggattt      60
atatatccat ccatccgcga cactttcccg ttgtattcta tcccttagtc ttcccttctc      120
cccctttcct tcttcacttc accttg ctc taa ccg aaa gta aac ctt tcc gcg      173
                        Leu      Pro Lys Val Asn Leu Ser Ala
                        1              5

atg aaa ttc tcc aat tct cta tcc gct ctc ctc gta tcc gca aat ctc      221
Met Lys Phe Ser Asn Ser Leu Ser Ala Leu Leu Val Ser Ala Asn Leu
      10              15              20

atg ttg gca gcg aag gcc tac aaa gga gat gcc acc ttt tat gat cct      269
Met Leu Ala Ala Lys Ala Tyr Lys Gly Asp Ala Thr Phe Tyr Asp Pro
      25              30              35              40

ggg ctg gga gct tgt ggc cat acg aat cag gct cat gaa ctt gtc gtt      317
Gly Leu Gly Ala Cys Gly His Thr Asn Gln Ala His Glu Leu Val Val
                        45              50              55

gcc ctt cca tca gcc aaa tac ggc agc gga gac cat tgt tcc aag cat      365
Ala Leu Pro Ser Ala Lys Tyr Gly Ser Gly Asp His Cys Ser Lys His
                        60              65              70

gtc ggc atc cac tac aaa ggc aaa tac gtg aaa gcc aaa gta gtc gac      413
Val Gly Ile His Tyr Lys Gly Lys Tyr Val Lys Ala Lys Val Val Asp
                        75              80              85

aaa tgt ccc ggt tgt ggt tcg aac gat tta gac atc tca cca acc gca      461
Lys Cys Pro Gly Cys Gly Ser Asn Asp Leu Asp Ile Ser Pro Thr Ala
                        90              95              100

ttc tct cac tta gcc agt caa gac ctc ggc cgt atc aaa gta gat tgg      509
Phe Ser His Leu Ala Ser Gln Asp Leu Gly Arg Ile Lys Val Asp Trp
      105              110              115              120

gaa ttt ctc tga tatcccatTT tcaatccctt acacgaaatc tgtatttgta      561
Glu Phe Leu

gaagaaagtc atgacgttat atagatcact tacatagatc ttcagggtttt cgtagatcga      621
cgaccgcgacgc tcttaaatat ttatttcccg tttttcygtt ttgtttttaa aaaaaaaaaa      681
aaaaaaaaa      689

```

<210> 7

<211> 122

<212> PRT

<213> Agaricus bisporus

<400> 7

Pro Lys Val Asn Leu Ser Ala Met Lys Phe Ser Asn Ser Leu Ser Ala
 1 5 10 15

Leu Leu Val Ser Ala Asn Leu Met Leu Ala Ala Lys Ala Tyr Lys Gly
 20 25 30

Asp Ala Thr Phe Tyr Asp Pro Gly Leu Gly Ala Cys Gly His Thr Asn
 35 40 45

Gln Ala His Glu Leu Val Val Ala Leu Pro Ser Ala Lys Tyr Gly Ser
 50 55 60

Gly Asp His Cys Ser Lys His Val Gly Ile His Tyr Lys Gly Lys Tyr
 65 70 75 80

Val Lys Ala Lys Val Val Asp Lys Cys Pro Gly Cys Gly Ser Asn Asp
 85 90 95

Leu Asp Ile Ser Pro Thr Ala Phe Ser His Leu Ala Ser Gln Asp Leu
 100 105 110

Gly Arg Ile Lys Val Asp Trp Glu Phe Leu
 115 120

<210> 8

<211> 2782

<212> DNA

<213> Agaricus bisporus

<220>

<221> misc_feature

<222> (2774)..(2774)

<223> n is an unknown nucleotide

<400> 8

ctgcgaattg gcataagcac ttraactttc gtcttcctca ctctcttcag gagattgaga 60
 ctgcatcggg gcaagcgagg gttgccgtac cgccctttga gaccccgaaa caagggattc 120
 gtcaagagca tcaagaagac tagaacgacc ggctgttttt ccacccgaca tcatagcaca 180
 aactgtcata aaccctgtgt caaaggggaa aaacaggcag agagaaggaa gggacgcgtc 240
 gcgatgaaat gtcattaac ctgaatgaca aacttcgcw aataacaatt taattaaaaa 300
 taaaaatcac gaggtgacaa acaggggtgt ttacctccat tcgactgcat cctggctctt 360

cccttccata gaactgtcgt ccatgcatgc accgctagca tcgcactgct ctgactcgca 420
taaccttaaa acgcggtggac cccctgttcg gacggccggt tcaggatccg gggctcagga 480
cacagtaaaa tcacaaaaac tcatactttg agagatatga cttctcgact tgcgccttcg 540
atggacggac aaattatccc caggtaccgg atctgtgaca ccgaattagt gcgcgatatt 600
atatatgact tttagcgggc gtctcatacg accgctcaag tccttgggga tggagaatgt 660
cacctcctgg tccaccgggc ccagagcatt acccggtcat taatctagcg cttcttgcatt 720
gcaactcctgc atgatcacc cagcgggcgg cgttttatcg gacatataag gaacaagatt 780
ccataggtag tggatcccct actccacctc cgcctactt ttataccaac cccaaatcca 840
aaggttgaaa aaaaaatttc gacaaggatt tatatatcca tccatccgcg acactttctc 900
gtttgattct atcccttagt ctttccttct ccccctttcc ttcttcaact cacttgctc 960
taaccgaaag taaaccttcc cgcgatgaaa ttctccaatt ctctatccgc tctcctcgta 1020
tccggaatc tcatgttggc agcgaaggcc tacaaaggag atggtcagtt atgacagtgt 1080
cgcccatcct atcgtcaatg atgctgattt tgttctccca ccttttttagc caccttttat 1140
gacctcgttg tacgtttttc ttacactttt caccaagatt cctcaccac tggtttcaaa 1200
attagctggg agcttgtggc catacgaatc aggtcatga acttgttgtt gcccttccat 1260
cagccaaata cggcaacgga gaccattgtt ccaagcatgt cggcatccac tgtacgtttc 1320
ctccttctcc ccacttctca aaaatcaaaa ttttactcat tgtaacaaga caaaggcaaa 1380
tacgtgaaag tcaaagtagt cgacaaatgt cccggctgtg gttcgaacga tctagacatc 1440
tcaccaaccg cattctctca gttagccagc caagacctcg gccgtatcaa agtagattgg 1500
gaatttatct aatatcccat ttcaatccct tacacgaaat ctgtacttgt agtcttagaa 1560
gaaagtcag acgttatata gatcacttac atagatcttc aggttttcgt agatcgacga 1620
ccgacgctct taaatattta ttacagttt ttctgtttt tgttttattg tcgcttggat 1680
ataagggtgt atactttgat atgattgcct acacacatat atcaacacag ttttagttat 1740
atcaacatca aaacatcagt caaggaaaac aaagagcgaa cgataaacat cagcacaagt 1800
atgtcagatt atggccaag aacgcgaaaa gaagttcgca aaagaacaga aactatcga 1860
aaagtgcaga tacataggtc acacaattaa cgacttccc gaatagttcc ctccaacctc 1920
ttatcgcgac tactagcacc aacggtaaca ccaaaagtac cttcaggcct cctccatccc 1980
tgtgcattca catcccaaat actcaaatca tacctcgaca aggtcatttt tacattccta 2040
gtctctccag ggccaatcgg tacagagtcg aaaccgcgta gcacggaagg aggttctcca 2100
gcagattcag ggaagttaat gtagagttgg ggagactcgg caccctaaaag tcgaccggta 2160
ttcttgacgt tgaaggaaac ctctgacaaa ggacgatgga gcctggaaac cggtcagcgt 2220
gaaattgttg rttagagaac cctcggcgct taccaagaag cgatagaacc accttcaact 2280
tgcgggcttg catgtccttg atcccatcc ctagccgcat caacttctct accttgacg 2340
tgcactttct taatagatat cttcgagtaa tcaaacttcg tgtaactcaa gccaaaacca 2400

aattcaaacc gaggtctaat gccattctgc aagagaaaat tcagtaagcg aacgtttccg 2460
agagaatcaa gcagcactgg cttacagcgt cgaagtgccg gtaatcgatt tgcaaacgat 2520
catcgtatgg tatagaaaga atatctcgaa taccatctcc ggggataact tgggccgaat 2580
aatcttcaat ccgtttggca atggtatatg gaagccttcc ggatgggttc caatctccat 2640
ataggacatc ggtcaacgaa tttcccgctt ctgtcccggt gcccagccc aaagaacctg 2700
gaaatagtca acaccggcgt tcacacaca agtagaagat acgaacagca gtgacattag 2760
gatgattgat ccanggttca ag 2782

<210> 9

<211> 984

<212> DNA

<213> *Agaricus bisporus*

<400> 9
ctgcgaattg gcataagcac ttraactttc gtcttcctca ctctcttcag gagattgaga 60
ctgcacgggt gcaagcgagg gttgccgtac cgccctttga gaccccgaaa caagggattc 120
gtcaagagca tcaagaagac tagaacgacc ggctgttttt ccacccgaca tcatagcaca 180
aactgtcata aaccctgttt caaaggggaa aaacaggcag agagaaggaa gggacgcgtc 240
gcatgaaat gctcattaac ctgaatgaca aacttcgcw aataacaatt taattaaaaa 300
taaaaatcac gaggtgacaa acaggggtgt ttacctccat tcgactgcat cctggctctt 360
cccttcata gaactgtcgt ccatgcatgc accgctagca tcgactgct ctgactcgca 420
taaccttaaa acgcgtggac cccctgttcg gacggccggt tcaggatccg gggctcagga 480
cacagtaaaa tcacaaaaac tcatactttg agagatatga cttctcgact tgcgccttcg 540
atggacggac aaattatccc caggtaccgg atctgtgaca ccgaattagt gcgcgatatt 600
atatatgact tttgacgggc gtctcatacg accgctcaag tccttgggga tggagaatgt 660
cacctcctgg tccaccgggc ccagagcatt acccggtcat taatctagcg cttcttgcatt 720
gcactcctgc atgatcacc caccggccg cgttttatcg gacatataag gaacaagatt 780
ccataggtag tggatcccct actccacctc cgcctactt ttataccaac cccaaatcca 840
aagggtgaaa aaaaaatttc gacaaggatt tatatatcca tccatccgcg acactttctc 900
gtttgattct atcccttagt ctttccttct cccctttcc ttcttcactt caccttgctc 960
taaccgaaaag taaacctttc cgcg 984

<210> 10

<211> 1270

<212> DNA

<213> Agaricus bisporus

<220>

<221> misc_feature

<222> (1262)..(1262)

<223> n is an unknown nucleotide

<400> 10
tatcccattt caatccctta cacgaaatct gtacttgtag tcttagaaga aagtcattgac 60
gttatataga tcacttacat agatcttcag gttttcgtag atcgacgacc gacgctctta 120
aatatatttatt tacagttttt ctgttttttg ttttattgtc gcttgatata aaggtggtat 180
actttgatata gattgcctac acacatatat caacacagtt ttagttatat caacatcaaa 240
acatcagtcac aggaaaacaa agagcgaacg ataaacatca gcacaagtat gtcagattat 300
ggccaagaa cgcgaaaaga agttcgcaaa agaacagaac actatcgaaa agtgcagata 360
cataggtcac acaattaacg acttcccga atagttccct ccaacctctt atcgcgacta 420
ctagcaccaa cggtaacacc aaaagtacct tcaggcctcc tccatccctg tgcattcaca 480
tcccaaatac tcaaatcata cctcgacaag gtcattttta cattcctagt ctctccaggg 540
ccaatcggta cagagtcgaa accgcgtagc acggaaggag gttctccagc agattcaggg 600
aagttaatgt agagttgggg agactcgga cccaaaagtc gaccggtatt cttgacgttg 660
aaggaaacct cgtacaaagg acgatggagc ctggaaaccg gtcagcgtga aattgttgrt 720
tagagaaccc tcggcgctta ccaagaagcg atagaaccac cttcaacttg cgggcttgca 780
tgtccttgat ccatttcctt agccgcatca acttctctac ctttgacgtg cactttctta 840
atagatatct tcgagtaatc aaacttcgtg taactcaagc caaaaccaa ttcaaaccga 900
ggctcaatgc cattctgcaa gagaaaattc agtaagcgaa cgtttccgag agaataaagc 960
agcactggct tacagcgtcg aagtgccggt aatcgatttg caaacgatca tcgtatggta 1020
tagaaagaat atctcgaata ccattctcgg ggataacttg ggccgaataa tcttcaatcc 1080
gtttggcaat ggtatatgga agccttccgg atgggttcca atctccatat aggacatcgg 1140
tcaacgaatt tcccgttct gtcccgtgac ccagcccaa agaacctgga aatagtcaac 1200
accggcgttc atcacacaag tagaagatac gaacagcagt gacattagga tgattgatcc 1260
anggttcaag 1270

<210> 11

<211> 835

<212> DNA

<213> Agaricus bisporus

<400> 11
 ccttgccgtt ttccagaagc tgcgaccagt cctcggagga gggggaacta aagatgtcga 60
 aagcagcagg gtgtgtgaca tagcgggaag aaagggatgat gatgatgaga acgaggggta 120
 cgatgaggac ggcccatctg aattgacggc caacgcgacg tttccggtcg tgagaggaca 180
 tggcaaagga gacgggggga ggggcgaggg tggcggagga ggtgctcgtg ccgaattcgg 240
 cacgagctca ccatgaaatt cgcaactgct ctccctgcct gccttactgc tgctgctagc 300
 gctcarcggc tcctcatcgg atccccctcct gaccaagcaa atctttctgc tggccagaac 360
 actacgattc aaattgtact cccgaatttc caatcgtctt cgcaagaagt tgcggtagtg 420
 cttgggatca cgtcctgcgc cgctgctccc tggcctgctc cagccgatac gatgggtcgt 480
 atcctttaca gcggtcattt caaccgcag agagatcctg caatgcccgc aatgcaagcc 540
 tacgaaaatt tcacggtctt cttgccggag aacctgccta agggcgcggc gcagattaac 600
 gtttaccatg tcgcacttat cggggccgggt ctcatgccat ggaacgagac gttgtccacc 660
 acagctttga ttcagtaatt ccatcaggat ttgaaatgga cctttagtag ttactgttt 720
 tgctatcgaa cgattcgrat aattacctga gatcaggctg gtgactgagg cccgtcggag 780
 tgctaccata atggcataat aaaattatac tcagctgaaa aaaaaaaaaa aaaaa 835

<210> 12

<211> 770

<212> DNA

<213> Agaricus bisporus

<400> 12
 tcgggttctc aagctacgat aaccgttcct cacgcatcac attcttctct gattgtaatg 60
 accgaagacc gcggttccac ccctaataatt cgaagagctt gttgccaatg tcaaaaaagg 120
 cttttgggca aaccgggatg gcttatcgag cctccagccc cagcagctaa catcgggcag 180
 agaaggaaaa tcacggcgtg tgaattatca ccgtttggtt cctgagtcac ctggagatgt 240
 acgcagatgg tgataccgtg tttgattggc gccgttgag aagaactata ttattcgatg 300
 gattttttgt tcgagtttga cacagagaca gagatgatag aggtttgcta ttgatgtagc 360
 aaaggatcat ttgacgatgg cgcatagggc gatggttacc tttatgtctg gaattataat 420
 atgtattggt ccccaactttt cttttatatt tattaatact aattggaagt ttcagttggt 480
 ggatgagcaa agttgggtgca gatagaaact agaattcgga ttcccatatc tgaggtaacct 540
 tttccttccg ctggcaatcc tggccacttc gacgtggtga cgcagagggc gcgtgctatt 600
 gtttagacat gccatatgga tcgacgttgc ctctcgtact tcgcgcctag gctcgtcat 660
 gcctcgatgc atctttcaat tcgggcgttg cgtctcccag gtgcctgtta aaagggcgaa 720
 ctttagtgta attgtactaa cacagtcctt cgggctgagc tctattcatc 770

<210> 13
<211> 703
<212> DNA
<213> *Agaricus bisporus*

<400> 13
ataacaattt aattaaaaat aaaaatcacg aggtgacaaa caggggtgtt tacctccatt 60
cgactgcac cttgctcttc ccttccatag aactgtcgtc catgcatgca ccgctagcat 120
cgcactgctc tgactcgcat aaccttaaaa cgcgtggacc ccctgttcgg acggccgggt 180
caggatccgg ggctcaggac acagtaaaat cacaaaaact catactttga gagatatgac 240
ttctcgactt gcgccttcga tggacggaca aattatcccc aggtaccgga tctgtgacac 300
cgaattagt cgcgatatta tatatgactt ttgacgggcg tctcatagca ccgctcaagt 360
ccttggggat ggagaatgtc acctcctggt ccaccgggcc cagagcatta cccggtcatt 420
aatctagcgc ttcttgcacg cactcctgca tgatcacccc acgcggccgc gttttatcgg 480
acataataag aacaagattc cataggtagt ggatccccta ctccacctcc cgcctacttt 540
tataccaacc ccaaatacaa aggttgaaaa aaaaatttcg acaaggattt atatatccat 600
ccatccgcga cactttctcg ttgattcta tcccttagtc tttccttctc cccctttcct 660
tcttcacttc accttgctct aaccgaaagt aaacctttcc gcg 703

<210> 14
<211> 486
<212> DNA
<213> *Agaricus bisporus*

<220>
<221> misc_feature
<222> (4)..(9)
<223> Restriction site for KpnI

<220>
<221> misc_feature
<222> (477)..(482)
<223> Restriction site for NarI

<220>

<221> Intron

<222> (277) .. (328)

<223>

<220>

<221> Intron

<222> (349) .. (408)

<223>

<220>

<221> Intron

<222> (415) .. (468)

<223>

<400> 14

cggggtaccg aggtccgcaa gtagattgaa agttcagtac gtttttaaca atagagcatt	60
ctcgaggctt gcgtcattct gtgtcaggct agcagtttat aagcgttgag gatctagagc	120
tgctgtttcc gcgtctcgaa tgttctcggg gtttaggggt tagcaatctg atatgataat	180
aatttgatgat gacatcgata gtacaaaaac cccaattccg gtcacatcca ccatctccgt	240
tttctcccat ctacacacaa caagcttata gccatgggtt gtctctcgct tgcataccat	300
ccagcagctc actgatgtcg acttgtagggt taaagttgga atcaacgggt aagtgttttt	360
gtcgtcgcgc tgtgggtccg gatcatctca gactttgggt gtcttgagcgt ttcggtgagt	420
gaccaccctg cattctggct atatgcgtga tactgacat cgctcaagggt cgtatcggcg	480
ccggcc	486

<210> 15

<211> 57

<212> DNA

<213> Agaricus bisporus

<220>

<221> CDS

<222> (1) .. (57)

<223>

<400> 15
atg cat ttc tct ttg tct ttt gcc acc ctt gct ctc tta gtc gct tcg 48
Met His Phe Ser Leu Ser Phe Ala Thr Leu Ala Leu Leu Val Ala Ser
1 5 10 15

gct gtt ggt 57
Ala Val Gly

<210> 16

<211> 19

<212> PRT

<213> Agaricus bisporus

<400> 16

Met His Phe Ser Leu Ser Phe Ala Thr Leu Ala Leu Leu Val Ala Ser
1 5 10 15

Ala Val Gly

<210> 17

<211> 45

<212> DNA

<213> Agaricus bisporus

<400> 17
gtgggattat ggtagccaa atggtcgtag ctaattatct tgcag 45

<210> 18

<211> 47

<212> DNA

<213> Agaricus bisporus

<400> 18
gtacgtaggc aagtggttcc caaaaagcac cagctaacac aacgtag 47

<210> 19

<211> 48

<212> DNA

<213> Agaricus bisporus

<400> 19
gtacgttgaa tcgtacaaga aagtgtaatc atcctgactt tctatcag 48

<210> 20

<211> 62

<212> DNA

<213> Agaricus bisporus

<400> 20
gtaagcaccg gctgcggttcg cacacccgtc ttgtgaaaag tcgtctcatg aatatcgccc 60
ag 62

<210> 21

<211> 51

<212> DNA

<213> Agaricus bisporus

<400> 21
gtaattaacc tcatcattat tgatccttcc catgcttaca gctgttatca g 51

<210> 22

<211> 52

<212> DNA

<213> Agaricus bisporus

<400> 22
gtgcggttctt ctcttgctct atcacaaatt ctgacgccgc aggacttgcc ag 52

<210> 23

<211> 63

<212> DNA

<213> Agaricus bisporus

<400> 23
gcaagcttct cttgtcacaa tgtaacggc gaggggtctg actcccttgg ttgtttttgt 60
tag 63

<210> 24

<211> 48

<212> DNA

<213> Agaricus bisporus

<400> 24

gtacgctata gcttgcaagg atggacatat ctaatcgggg acgtgtag

48

<210> 25

<211> 61

<212> DNA

<213> Agaricus bisporus

<400> 25

gtgggttcca tcatctcttg agttatgccg cggtgactg atcatgtttc taatacttca

60

g

61

<210> 26

<211> 46

<212> DNA

<213> Agaricus bisporus

<400> 26

gtgcgtcatg tccgtatcat ctactctcat actaatgcgc atatatg

46

<210> 27

<211> 53

<212> DNA

<213> Agaricus bisporus

<400> 27

gtagggccct tttatctttg gtccgattgc gtcattcatg tctatatatg cag

53

<210> 28

<211> 48

<212> DNA

<213> Agaricus bisporus

<400> 28

gtagggttta ttcacccctc tggtcttata agcttgacat ccttcaag

48

<210> 29

<211> 53

<212> DNA

<213> Agaricus bisporus

<400> 29

gtaagcataa ggagttgctg caggcgaccc atagcgttct aattagctgc tag

53

<210> 30

<211> 52

<212> DNA

<213> Agaricus bisporus

<400> 30

gtatgttgga actcaccgat gcgctcttgg ttgattttat tttctacat ag

52

<210> 31

<211> 50

<212> DNA

<213> Agaricus bisporus

<400> 31

gtgggtgac attcgagggt gtcttctctgt gtattgataa ggtttgctag

50

<210> 32

<211> 45

<212> DNA

<213> Agaricus bisporus

<400> 32

gtcagtacta gtttgtttct tcttacacct tctcattctt tgcag

45

<210> 33

<211> 56

<212> DNA

<213> Agaricus bisporus

<400> 33
gtgagcactg caatatggta tagcttggaa agcctttatt tatacagaac atccag 56

<210> 34

<211> 55

<212> DNA

<213> Agaricus bisporus

<400> 34
gtgagtgaag atagtcttca tgtgagatcc ttgtactaat tgccgaacgt cgtag 55

<210> 35

<211> 690

<212> DNA

<213> Agaricus bisporus

<400> 35
tatatagtct ccaaattcta ttgtaatgcc attttcccaa ttcaaaagga cccgctctcg 60
aaccgggtca gatgcaattt tggtcagcaa tggtttatgt tgtttcccg taagtatgca 120
ctagaagaga acaaaacgtc actatttgct caatgcagga tgcacctggc gagataatat 180
tcttgcggtg aagtcgaaca acgtctgtag tcctgtaaaa atatacagtg agtagaggga 240
tgatgccgat gtggaaggaa gcaaccgatt acgattcggg tttggccatg agacggccgc 300
tcttgaacag accaacaata tccctttaaa tttaatacag aattactcaa tatgcttcca 360
agtatttcga ttcctcgaat tccgtctggc cgcggcatga gcatggacag gcgacagaa 420
gaggctatcg ttgtattgct tcatcagcga ccctgactag tgacttcagg catgatcatg 480
cgcttagcaa tctgtccctt caagtcgagt ccccgaaattc aacagcttca acaagtcgtg 540
attatttgac ccccgactgg aatcaaattg gctcttcaaa tttcaaactt caatgcttca 600
tgcttcatgc gtcatgacgc aagctgtcaa ttttcatttt ccagttcggg cccattotca 660
ctctcgcccc tcctaattgc ttccagaaaa 690

<210> 36

<211> 910

<212> DNA

<213> Agaricus bisporus

<400> 36
tatcccatth caatccctta cacgaaatct gtactttag tagcttagaaga aagtcatgac 60
gttatataga tcacttacat agatcttcag gttttcgtag atcgacgacc gacgctctta 120

aatatatttatt tacagttttt ctgttttttg ttttattgtc gcttgatat aagggtggtat 180
actttgatat gattgcctac acacatatat caacacagtt ttagttatat caacatcaaa 240
acatcagtcac aggaaaacaa agagcgaacg ataaacatca gcacaagtat gtcagattat 300
ggtccaagaa cgcgaaaaga agttcgcaaa agaacagAAC actatcgaaa agtgcagata 360
catagggtcac acaattaacg acttcccga atagtccct ccaacctctt atcgcgacta 420
ctagcaccaa cggtaacacc aaaagtacct tcaggcctcc tccatccctg tgcattcaca 480
tcccaaatac tcaaatcata cctcgacaag gtcattttta cattcctagt ctctccaggg 540
ccaatcggtc cagagtcgaa accgcgtagc acggaaggag gttctccagc agattcaggg 600
aagttaatgt agagttgggg agactcggca cccaaaagtc gaccggtatt cttgacgttg 660
aaggaaacct cgtacaaagg acgatggagc ctggaaaccg gtcagcgtga aattgttgat 720
tagagaaccc tcggcgctta ccaagaagcg atagaaccac cttcaacttg cgggcttgca 780
tgtccttgat cccattccct agccgcatca acttctctac ctttgacgtg cactttctta 840
atagatatct tcgagtaatc aaacttcgtg taactcaagc caaaaccaa ttcaaaccga 900
ggctcaatgc 910

<210> 37

<211> 800

<212> DNA

<213> *Agaricus bisporus*

<220>

<221> misc_feature

<222> (5)..(9)

<223> Restriction site for BglII

<220>

<221> misc_feature

<222> (790)..(795)

<223> Restriction site for KpnI

<220>

<221> misc_feature

<222> (290)..(290)

<223> n is an unknown nucleotide

<220>

<221> misc_feature

<222> (498)..(498)

<223> n is an unknown nucleotide

<220>

<221> misc_feature

<222> (589)..(589)

<223> n is an unknown nucleotide

<400> 37

gaatagatct gatgccgacc gcgggatcca cttaacgtta ctgaaatcat caaacagctt	60
gacgaatctg gatataagat cgttggtgtc gatgtcagct ccggagttag gacaaatggt	120
gttcaggatc tcgataagat acgttcattt gtccaagcag caaagagtgc cttctagtga	180
tttaatagct ccatgtcaac aagaataaaa cgcgttttcg ggtttacctc ttccagatac	240
agctcatctg caatgcatta atgcattgac tgcaacctag taacgccttn caggctccgg	300
cgaagagaag aatagcttag cagagctatt ttcatatttc ggagacgaga tcaagcagat	360
caacggtcgt caagagacct acgagactga ggaatccgct cttggctcca cgcgactata	420
tatttgcttc taattgtact ttgacatgct cctcttcttt actctgatag cttgactatg	480
aaaattccgt caccagcncc tgggttcgca aagataattg catgtttctt ctttgaactc	540
tcaagcctac aggacacaca ttcacgttag gtataaacct cgaaatcant tcctactaag	600
atggtataca atagtaacca tgcattggtg cctagtgaat gctccgtaac acccaatacg	660
ccggccgaaa cttttttaca actctcctat gagtcgttta ccagaaatgc acaggtacac	720
ttgttttagag gtaatccttc tttctagaag tcctcgtgta ctgtgtaagc gccactcca	780
catctccacg gtacctgcag	800